

ELECTRON EXCITATION CROSS SECTIONS FOR THE S // TRANSITIONS $3s^23p^3\ ^4S^\circ \rightarrow 3s^23p^3\ ^2D^\circ, ^2P^\circ$ and $3s3p^4\ ^4P$

C. Liao¹, Steven J. Smith¹, D. Hitz², A. Chutjian¹, and S. S. Tayal³

¹Jet Propulsion Laboratory/Caltech, Pasadena, CA 91109 USA

²CEA/DRFMC, 38054 Grenoble Cedex 9, France

³Department of Physics, Clark Atlanta University, Atlanta, GA 30314 USA

Emission lines from excited S // are regularly detected in a wide variety of astronomical objects, including the Io torus, Sun, planetary nebulae, H II regions, Seyfert galaxies, and supernova remnants. Experimental and theoretical collisional excitation cross sections are reported here for the observed transitions $3s^23p^3\ ^4S^\circ \rightarrow 3s^23p^3\ ^2D^\circ, ^2P^\circ$ and $3s3p^4\ ^4P$ in S //. The transition wavelengths (energies) are 6716 Å (1.85 eV), 4069 Å (3.05 eV), and 1256 Å (9.87 eV), respectively. In the experiments, use is made of the energy-loss, merged-beams method^{1,2}. The metastable fraction of the S // beam is assessed and minimized. The contribution of elastically-scattered electrons is reduced by the use of lowered solenoidal magnetic field, use of a modulated radiofrequency voltage on the analyzing plates, and retarding grids to reject the elastically-scattered electrons with larger Larmor radii. For each transition, comparison is made between present experiment and 19-state R-matrix calculation, and with three previous close-coupling calculations^{3,4,5}.

Shown in Fig. 1 for the lowest-lying, forbidden $^4S^\circ \rightarrow ^2D^\circ$ transition are present experimental results, with comparisons to four theoretical calculations: the present 19-state close-coupling (CC) calculation, a 6-state CC calculation³, a 12-state CC calculation⁴, and an 18-state CC calculation⁵. One finds general excellent agreement among the experiment and theories, from threshold to 6 eV (experiment-theories) and 10 eV (theories alone).

In Fig. 2 are presented results for the second forbidden $^4S^\circ \rightarrow ^2P^\circ$ transition. Here one finds considerable discrepancy in the location and cross section of the peak. Experiment favors results in the 18-state and 19-state CC theories, while the 12-state and 6-state theories differ in shape, and give about one-half the experimental cross section.

C. L. and D. H. thank the NAS-NRC for support at JPL. S. S. T. acknowledges support from the NASA under the Planetary Atmospheres Program. The experimental work was carried out at JPL/Caltech, and was supported by NASA.

References

1. S. J. Smith, A. Chutjian, J. Mitroy, R. Henry, K-F. Man, R. Mawhorter, and I. Williams, *Phys. Rev. A* **48**, 292 (1993); S. J. Smith, M. Zuo, A. Chutjian, S. Tayal, and I. Williams, *Astrophys. J.* **463**, 808 (1996).

2. M. E. Bannister, X. G. Guo, T. M. Kojima, and G. H. Dunn, *Phys. Rev. Letters* **72**, 3336 (1994).

3. S. S. Tayal, R. J. W. Henry, and S. Nakazaki, *Astrophys. J.* **313**, 487 (1987).

4. W. Cai and A. K. Pradhan, *Astrophys. J. Suppl. Ser.* **88**, 329 (1993).

5. C. A. Ramsbottom, K. L. Bell, and R. P. Stafford, *At. Data Nucl. Data Tables*, **63**, 57 (1996).




Figure 1. Experimental (filled circles) and theoretical cross sections for excitation of the $^4S^\circ \rightarrow ^2D^\circ$ transition. Theoretical results are convoluted with a 250 meV (FWHM) electron-energy width: solid line, present 19-state R-matrix calculation; dotted line, Ref. 3; dashed line, Ref. 4; long-short dashed line, Ref. 5.




Figure 2. Experimental and theoretical cross sections for excitation of the $^4S^\circ \rightarrow ^2P^\circ$ transition. Theoretical results are convoluted with a 250 meV (FWHM) electron-energy width. Notation for the theories is the same as in Fig. 1.